

GALAXY LUMINOSITY FUNCTION IN GROUPS

HERNÁN MURIEL, CARLOS A. VALOTTO AND DIEGO G. LAMBAS

Grupo de Investigaciones en Astronomía Teórica y Experimental (IATE)

Observatorio Astronómico de Córdoba, Laprida 854, 5000 Córdoba, Argentina

CONICET, Buenos Aires, Argentina

hernan@oac.uncor.edu, val@oac.uncor.edu, dgl@oac.uncor.edu

Draft version February 1, 2008

ABSTRACT

We compute the luminosity function of galaxies in groups. The data consists in two samples of groups of galaxies selected in distance and redshift space comprising a total number of 66 groups. The assignment of galaxies to the groups were derived from the Edinburgh-Durham Southern Galaxy Catalog, considering a limiting apparent magnitude $m_{lim} = 19.4$.

We find a flat faint end of the mean galaxy luminosity function in groups in contrast to the mean galaxy LF in clusters where a large relative number of faint galaxies is present. We find that a Schechter function with parameters $M^* \simeq -19.6 \pm 0.2$, $\alpha \simeq -1.0 \pm 0.2$ provides a satisfactory fit to the galaxy LF of our total sample of groups in the range of absolute magnitudes $-22 < M < -16$.

1. INTRODUCTION

Several works have attempted the identification of groups of galaxies in 2 dimensions using catalogs with angular positions (see for instance Turner and Gott 1976). A major problem with such identifications is the low density contrast of these systems and therefore redshift surveys are essential to provide reliable physical groups. Nevertheless, most of redshift surveys are limited to bright apparent magnitudes allowing only the identification of the brightest members. This fact make difficult the determinations of the faint end of the luminosity function (hereafter LF) of galaxies in groups. The galaxy LF in clusters and nearby groups has been studied by several authors (Oemler 1974, Dressler 1978, Oegerle et al. 1986, Oegerle et al. 1987, Gudehus and Hegyi 1991, Willmer et al. 1990, Garilli et al. 1991, Ferguson and Sandage 1991, Garilli et al. 1992, Barrientos et al. 1996, Gaidos 1997, Valotto et al. 1997, Trentham 1997). Of particular importance is the possible universal character of the galaxy LF early suggested by Abell 1962 and Abell 1975 since it may serve to pose constraints to models of galaxy formation as well as testing the importance of environment in the related astrophysical processes. Interactions play a fundamental role in galaxy evolution (see for instance Postman and Geller 1984) being particularly relevant in small galaxy associations. Moreover, in contrast to clusters, groups have a low galaxy velocity dispersion, gas density and temperature which may cause a significantly different evolution of galaxies in groups and clusters. Groups of galaxies are therefore suitable systems to provide a useful observational insight on different physical phenomena related to environment.

The field galaxy LF has been determined in several works, see for instance Loveday et al. 1992, and more recently Marzke et al. 1994 and Lin et al. 1996. These authors provide Schechter function fits (Schechter 1976) and in spite of discrepancies in the value of the parameter M^* , both determinations are consistent with a flat LF at the faint end, $\alpha \simeq -1$. On the other hand, several studies of the galaxy luminosity function in clusters show a much

steeper galaxy LF at the faint end, $\alpha \leq -1.4$ (e.g. Ferguson and Sandage 1991, Valotto et al. 1997, Trentham 1997).

The universality of the galaxy LF in clusters has been seriously questioned by López-Cruz & Yee 1995 and López-Cruz et al. 1997. From their sample of 45 Abell clusters with $z < 0.14$ 39 clusters show an increase of the fraction of dwarf galaxies consistent with a double Schechter fit with $\alpha \simeq 1$ and $-2 < \alpha < -1.4$ and only 7 can be suitably fitted with a single Schechter fit with $\alpha \simeq 1$. This set of clusters with flat LF are characterized by an evolved morphology of the cD type and are on average massive and gas rich. Nevertheless, Driver et al. 1995 and Trentham 1997 find rising LF in rich clusters which suggest the complexity of this phenomenon. In spite of the reported differences of the galaxy LF in clusters the results show agreement with the mean galaxy LF in clusters derived by Valotto et al. 1997 which rises at faint magnitudes consistent with a single Schechter fit with $\alpha \simeq -1.4$. The tendency of poorer clusters to show a flatter galaxy LF ($\alpha \simeq -1.2$) than richer clusters ($\alpha \simeq -1.5$) was obtained by Valotto et al. 1997 dividing their sample in two equal number of clusters. These results and López-Cruz et al. 1997 findings would not be necessarily inconsistent given that the flat LF clusters ($\alpha \simeq -1.0$) observed correspond to a small fraction ($\simeq 20\%$) of dynamically relaxed systems where evolutionary process have an important role. Other works provide useful studies of the galaxy LF in different environments. For instance, Gaidos 1997 develop an important work on the luminosity function of galaxies in clusters. Nevertheless, the background subtraction in this work is taken at 0.5 Mpc of the cluster center which may contribute significantly to the resulting flat LF of the analysis given the expected large contamination of the background by cluster members.

Given the observed differences between cluster and field mean galaxy luminosity functions we analyze in this work the galaxy LF in moderate density associations. In order to provide a suitable study of the effects of environment we use the same statistical procedures as in Valotto et al.

1997 avoiding possible systematic effects due to the use of different techniques. In section 2 we describe the group and galaxy data used. Section 3 gives a description of the methods of analysis adopted, the main results obtained and error estimates through a Monte-Carlo method. In Section 4 we present the main conclusions.

2. DATA

Willick et al. (1997) have assembled a homogeneous catalog of peculiar velocity data (the Mark III catalog). This catalog includes a sample of groups selected using redshifts and distances of the Mark III galaxies with a grouping algorithm as described in Willick et al. 1996. Although the authors claim that the dynamical characteristics of the groups are in any sense well defined, this sample is suitable for our analysis since our aim is to study the galaxy LF in moderate galaxy overdensities. In addition, completeness in group number is not required given that COSMOS galaxies are used to provide a statistical assignment to groups. The Mark III group sample comprise a total number of 277 objects. This sample of groups of galaxies has redshift independent distance estimates which enables us to study the galaxy LF for nearby groups of galaxies for which peculiar motions cannot be neglected.

Other samples of groups of galaxies without redshift independent distance estimates was taken from different group catalogs (Maia et al. 1989 (MCL), Fouqué et al. 1992 (FGCP) and Garcia 1993). García (1993) identify groups in the magnitude-limited (B_0 brighter than 14.0) sample of galaxies extracted from the Lyon-Meudon Extragalactic Database (LEDA). This author identify groups applying two different technique: the hierarchical clustering method (Materne 1978) and the percolation method (Huchra and Geller 1982). García identify 485 groups of at least three members and detected by both technique. MCL select groups of galaxies using the same technique applied by HG in the CfA catalog. This technique is apply to the Southern Sky Redshift Survey (da Costa et al. 1988). MCL identify 87 groups with more than two members and mean velocities smaller than 8000 km s^{-1} . FGCP list 246 groups of at least three members obtained by applying the hierarchical algorithm to the Catalog of Principal Galaxies (Paturel et al. 1989a, Paturel et al. 1989b). We have also included compact groups taken from Hickson (1982).

We have restricted this compilation of groups without redshift independent distance estimates to radial velocities $cz > 2500 \text{ km s}^{-1}$. With this restriction the group peculiar velocities would not seriously affect the estimates of absolute magnitudes due to peculiar motions. Since we aim to study moderate galaxy density enhancements we have not considered groups within $0.5 \text{ Mpc } h^{-1}$ in projection to Abell clusters of galaxies. We have adopted a limiting absolute magnitudes $M_{lim} = -16$ since the signal-to-noise ratio for fainter absolute magnitudes results too weak in our studies which impose a further restriction to group distances.

Our final sample comprises 66 groups, 35 with distances derived from the MarkIII catalog and 31 from our compilation of groups with redshift determinations from the literature from which 3 are Hickson compact groups. In figure 1 are displayed the distribution of distances for the two subsamples of groups.




Figure 1. Distribution of distances in km s^{-1} corresponding to 35 MarkIII groups and 31 groups with radial velocities (filled histogram) considered in our study.

The Edinburgh-Durham Southern Galaxy Catalog, hereafter COSMOS Survey (Heydon-Dumbleton et al. 1989), was used for the statistical assignment of galaxies to the groups. This survey provides angular positions and photographic magnitudes in the b_j band for over two million galaxies. We have restricted to the region $\delta < -10^\circ$ given the lower quality in the photographic material in the northern hemisphere. Valotto et al. (1997) suggest a limiting apparent magnitude $m = 19.4$ for statistical analysis in COSMOS Survey which minimizes errors due to misclassification of stars, galaxies plate variations, etc. The rms difference between CCD and COSMOS magnitudes is $\sigma \approx 0.25 \text{ mag}$ (Valotto et al. 1997, Loveday et al. 1992, Rousseau et al. 1996). This rms value has been used in the Monte-Carlo estimates of errors described in section 4.

TABLE 1 - Group Properties

RA [°]	Dec [°]	N_{gal}	$d [km \text{ s}^{-1}]$	Reference
2.7312	-33.5381		6488	MarkIII
9.2596	-27.3823		6427	MarkIII
9.3742	-20.1418	3	3474	MCL85
9.4075	-46.5883		3109	MarkIII
13.8497	-45.3011		6173	MarkIII
14.1764	-52.9298	3	7284	MCL19
17.1555	-46.2836	5	6621	MCL33
17.3560	-35.8672	3	6255	MCL54
18.0305	-31.7627		5951	MarkIII
18.5607	-33.2295		6011	MarkIII
18.6351	-32.1191	7	5345	MCL61
19.8244	-44.2044	4	6614	MCL36
19.9841	-33.9223		3187	MarkIII
20.1977	-44.8500		7091	MarkIII

TABLE 1 - Continued

21.6424	-23.2313	3	7350	HC11
22.1155	-35.7583	3	5308	MCL55
23.8488	-39.2949	3	5309	MCL47
24.4133	-27.4186		5713	MarkIII
24.8495	-42.7212		6012	MarkIII
26.9436	-35.0977		4715	MarkIII
29.2886	-50.4484		5546	MarkIII
31.3328	-55.1617	3	5725	MCL17
32.4185	-41.6438	3	5254	G50
32.3422	-22.5234		3538	MarkIII
32.8671	-23.2635		4668	MarkIII
33.0409	-39.4327	3	4924	MCL46
33.2765	-32.0978		3829	MarkIII
37.3372	-31.4779	3	4393	MCL64
37.4580	-42.9617	3	5027	MCL41
40.8292	-55.2613	3	5398	G74
41.3068	-17.6733	3	7232	MCL87
41.6544	-24.9411	3	6416	MCL75
41.8712	-55.3755	3	5483	MCL16
42.5540	-31.2784		5137	MarkIII
43.2061	-32.1921		5597	MarkIII
44.1086	-18.5158		3577	MarkIII
50.4756	-13.6459	7	9510	HC26
50.5104	-24.7825		5243	MarkIII
52.0631	-21.2036		2179	MarkIII
59.2783	-25.3789		3782	MarkIII
64.1942	-44.2942		4552	MarkIII
321.1159	-40.4012	4	4983	MCL45
322.5890	-39.5620	18	5145	G445
324.9505	-39.9815		4785	MarkIII
328.1029	-28.8809		6449	MarkIII
329.1513	-34.7459		2625	FGCP244
330.6722	-34.3896	3	4479	G451
330.8851	-32.1050		2842	MarkIII
331.3571	-18.9131		2749	MarkIII
331.5865	-27.8394	3	6770	MCL69
332.3014	-27.7760	4	7140	HC91
332.3542	-32.8460		3760	MarkIII
333.2208	-30.0362	3	4352	G454
333.7056	-27.4679	3	5190	MCL70
335.2614	-21.8241		5912	MarkIII
335.4913	-25.5694	7	4805	G457
336.5680	-28.2023		6551	MarkIII
336.5773	-14.9864	5	5066	G458
337.9965	-27.0941		3435	MarkIII
338.1969	-19.9765		7474	MarkIII
341.4786	-22.9033	3	3039	MCL76
341.5522	-39.8120		2555	MarkIII
341.7496	-22.6184		3369	FGCP251
342.9665	-20.1329		3184	MarkIII
351.5385	-35.3763		3177	MarkIII
352.9783	-46.7819		3267	MarkIII
354.5960	-47.6564	3	2781	MCL30

Table 1 lists the corresponding angular positions (J2000.0), published number of members N_{gal} when available and distances d (redshift independent estimates for MarkIII groups, and $d=c z$ otherwise) in km/sec. FGCP: Fouque et al., 1992; G: Garcia 1993; HC: Hickson 1982; MCL: Maia et al., 1989; MarkIII: Willick et al., 1997.

3. ANALYSIS AND RESULTS

3.1. Galaxy Counts and Background Subtraction.

We compute the number of galaxies brighter than a limiting absolute magnitude M_{lim} within a projected radial distance r from the centers of the groups. We have applied a K -correction term of the form $K = 2z$ (Esfthathiou et al. 1988). The group projected radius r was fixed at $0.5 h^{-1} Mpc$ where Hubble constant is $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$ similar to the adopted group radius in other studies (see for instance Geller and Huchra 1983, Garcia 1993).

We define a mean local background around each group in order to decontaminate the galaxy counts. This mean local background is defined as the number density of galaxies in the same range of apparent magnitudes in a ring at projected radii $R_1 < r < R_2$. According to Valotto et al. 1997 the stability of the results does not depend crucially on the adopted radius for background correction provided that the decontamination ring is well beyond the average projected radius of groups and small enough in order to take into account local variations of the projected galaxy density due to patchy galactic obscuration, large scale gradients in the galaxy catalog, etc. The counts of galaxies for each group are binned in magnitude intervals of 0.5 mag which is larger than the photometric errors. We subtract the corresponding mean background correction to each magnitude bin to compute the contribution from each group to the LF. The total decontaminated number of galaxies in the range $-22 < B_j < -16$ is $N_{tot} = 1280$ corresponding to an average number of galaxies $n \simeq 20$ per group which may be compared to $n \simeq 100$ in clusters (see Valotto et al. 1997, Table 2). We notice that in spite of this large difference in the number of galaxies, groups and clusters have comparable mean galaxy densities due to the radii adopted (0.5 and $1.5 h^{-1} \text{ Mpc}$ respectively).

3.2. Results and Error Estimates

Figure 2 shows the galaxy group LF of the total sample of groups with $M_{lim} = -16$. Error bars in this figure correspond to the Monte-Carlo determination of errors discussed below which provide a reliable estimates of the uncertainties.

Errors in the determination of the galaxy LF without individual galaxy distances arises from uncertainties in the decontamination of foreground and background galaxies. Valotto et al. 1997 found a good agreement between the galaxy cluster LF determinations from different background corrections although fluctuations in the background counts contribute significantly to errors. We have also considered the propagation of photometric errors in COSMOS magnitudes in order to provide suitable error estimates of the galaxy LF in groups. We use a Monte-Carlo algorithm that takes into account the observed scatter between CCD and COSMOS magnitudes through random Gaussian errors with dispersion $\sigma = 0.26 \text{ mag}$. added to galaxy magnitudes as well as errors associated to the different assignment of background corrections. For each group we obtain different background corrections by counting galaxies in circles of projected radius $r = 0.5 h^{-1} \text{ Mpc}$ centered at distances $2 h^{-1} \text{ Mpc} < r_1 < 4 h^{-1} \text{ Mpc}$ from the groups. We calculate the resulting LF in 50 random realizations which include magnitude errors and background fluctuations. Our Monte-Carlo estimates of errors in figure 2 correspond to the rms values of the relative number

frequency of galaxies in each absolute magnitude interval $M - M + \Delta M$.




Figure 2. Uncorrected luminosity function of galaxies in groups. Solid line corresponds to the Schechter best fit with parameters $M^* = -19.9$ and $\alpha = -1.0$. Dashed line correspond to the

In order to provide a suitable LF fit, we have adopted a Schechter function model $\phi(L)dL = \text{const} \times (L/L^*)^\alpha e^{-L/L^*} d(L/L^*)$ (Schechter 1976). We have applied a maximum-likelihood estimator using a χ^2 -minimization procedure developed by Levenberg and Marquard, (see Press et al. 1987) which minimizes the difference

$$\chi^2 \equiv \sum_{i=1}^N \left(\frac{\phi_i - \phi(L_i; \text{const}, \alpha, L^*)}{\sigma_i} \right)^2$$

where ϕ_i is the relative frequency of galaxies corresponding to the i^{th} luminosity bin and σ_i is its associated uncertainty. The problem is reduced to the derivation of the three parameters const , L^* and α which minimizes χ^2 . This method deals with errors giving a reliable set of fitting parameters provided the errors are representative of the true uncertainties involved. In our calculations we have considered errors derived through the Monte-Carlo estimates described above which we argue, give confident estimates of errors. In figure 2 is shown (solid line) the Schechter fit derived by the maximum-likelihood estimator with parameters $\alpha = -1.0$, $M^* = -19.9$ where can be seen a very good agreement between observations and the Schechter fit.

We have considered the effects on the Schechter function fitting parameters due to magnitude binning and photometric errors through a Monte-Carlo technique. We simulate a Schechter function LF with parameters corresponding to the group galaxy LF ($\alpha = -1.0$, $M^* = -19.9$) and the cluster galaxy LF ($\alpha = -1.4$, $M^* = -20.0$) provided by Valotto et al. 1997. For both cases we have considered

Gaussian photometric errors with dispersion $\sigma 0.25$ mag) and we have taken into account the adopted binning (0.5 mag, and 0.3 mag in groups and clusters respectively). In both cases we find a resulting M^* parameter $\simeq 0.3$ mag lower than the input values while the parameter α remains unchanged. The corresponding corrected fitting parameters for the group and cluster galaxy LF are $\alpha = -1.0$, $M^* = -19.6$) and $\alpha = -1.4$, $M^* = -19.7$ respectively.

A summary of the results obtained is given in Table 2. For comparison are also listed in this table the resulting Schechter fitting parameters of the corrected field galaxy LF (Loveday et al. 1992). It can be seen in this table the importance of environment on the shape of the LF. The group galaxy LF has a flat faint end consistent with the field galaxy LF in contrast to clusters where an important rise in the number of low luminosity galaxies is present.

TABLE 2 - Summary of Statistical Results

Samples	α	M^*
Groups	-1.0 ± 0.2	-19.6 ± 0.2
Rich Clusters	-1.5 ± 0.1	-19.7 ± 0.1
Poor Clusters	-1.2 ± 0.1	-19.6 ± 0.1
Field Galaxies	-1.0 ± 0.15	-19.5 ± 0.1

We show in figure 3 the corresponding 1σ and 2σ level error contours of corrected galaxy LF in groups (contours $\chi^2 - \chi^2_{ML}$) in the $\alpha - M^*$ plane. In this figure the corrected cluster and field galaxy LF fitting parameters given by Valotto et al. 1997 and Loveday et al. 1992 are also displayed.




Figure 3. Error contours 1σ (solid line) and 2σ (dashed line) of the Schechter function parameters α and M^* for the corrected galaxy LF in groups. The open circle and square correspond to the corrected cluster and field galaxy LF (Valotto et al. 1997 and Loveday et al. 1995).

4. CONCLUSIONS

We have applied statistical analyses to calculate the mean galaxy luminosity function in groups. We have used COSMOS survey galaxies and group catalogs with either available distances or redshifts. Schechter functions provide good fits to the galaxy LF of the sample of groups analyzed. The best fitting parameters for our total sample of 66 groups are $\alpha = -1.0 \pm 0.2$, $M^* = -19.6 \pm 0.2$. This value of the α parameter contrasts with that corresponding to a single Schechter fit in clusters $\alpha \approx -1.4$ (see for instance Ferguson and Sandage 1991, Valotto et al. 1997) indicating an important relative excess of faint galaxies in clusters. The dependence of the galaxy LF on cluster richness was explored by Valotto et al. 1997 finding that poor clusters show a flatter galaxy LF than more massive systems ($\alpha \simeq -1.2$ in contrast to $\alpha \simeq -1.5$). López-Cruz et al. 1997 results show a flat galaxy LF ($\alpha \simeq -1.0$) in a subsample of dynamically relaxed systems which may be a strong indication that evolutionary process may disrupt dwarf galaxies contributing to the formation of cD haloes in these systems. However, Driver et al. 1995 in their analyses of the galaxy LF of the rich (R=3) cD-dominated cluster A963 a large excess of faint galaxies is found which indicates the non universality of this phenomenon. Similar large excess of faint galaxies are reported by Trentham 1997 for 4 rich clusters (R=2-4) indicating that the dependence of the faint excess on different physical parameters

of clusters deserve further analysis. The importance of the cluster merging history and the gas-dynamical evolution within these systems should be addressed before firm conclusions can be derived from the analysis of the observations.

Our comparison with Valotto et al. 1997 results have the advantage that both have used similar methods of analysis, same galaxy catalog and background subtraction procedure and therefore it provides a reliable confrontation of the mean galaxy LF in clusters and groups. In spite of the fact that our sample galaxies correspond to moderate galaxy overdensities we find no evidence for an excess of faint galaxies relative to the field as it is found for the mean galaxy LF in clusters. Although galaxy mutual interactions are expected to play an important role in groups our results indicate that at these moderate galaxy density enhancements the galaxy luminosity function is not significantly different from the field.

We acknowledge Dr. MacGillivray for kindly permitting the use of the COSMOS Survey. We thank the Referee for useful comments and suggestions which greatly improved the original version of this paper. This work was partially supported by the Consejo de Investigaciones Científicas y Técnicas de la República Argentina, CONICET, the Consejo de Investigaciones Científicas y Tecnológicas de la Provincia de Córdoba, CONICOR, and Fundación Antorchas, Argentina.

REFERENCES

- Abell, G.O. 1962, in Problems in Extragalactic Research, IAU Symposium N^o15. Ed. G.C.McVittie (Mac Millan, New York), 213.
 Abell, G.O. 1975, in Star and Stellar Systems, Vol. 9. Ed. A. Sandage, M.Sandage & J.Kristian (University of Chicago, Chicago).
 Abell, G.O., Corwin, H.G. & Olowin, R.P. 1989, ApJS, 70, 1.
 Barrientos, L.F., Schade, D. & López-Cruz, O. 1996, ApJ, 460, L89.
 da Costa, L.N., Pellegrini, P.S., Sargent, W.L., Tonry, J., Davis, M., Meiksin, A., Latham, D.W., Menzies, J.W. and Coulson, I.A. 1988. ApJ, 327,544.
 Dressler, A. 1978, ApJ, 226,55.
 Driver, S.P., Philipps, S., Davies, J.I., Morgan, I. & Disney, M.J. 1994, MNRAS, 268,393.
 Efsthathiou, G., Ellis, R.S. & Peterson, B.A. 1988, MNRAS, 232,431.
 Ferguson, H.C., & Sandage, A. 1991, AJ, 101,765
 Fouque, P., Gourgoulhon, E., Chamaraux, P. and Paturel, G. 1992. A&A Suppl. Ser. 93,211.
 Gaidos, E., AJ, 113, 117.
 Garcia, A.M. 1993. A&A Suppl. Ser. 100,47.
 Garilli, B., Maccagni, D., & Vettolani, G. 1991, AJ, 101,795.
 Garilli, B., Bottini, D., Maccagni, D., Vettolani, G., & Maccacaro, T. 1992, AJ, 104,1290.
 Geller, M.J. and Huchra, J.P. 1983. ApJS52,61.
 Gudehus, D.H., & Hegyi, D.J. 1991, AJ, 101,18.
 Heydon-Dumbleton, N.H., Collins, C.A. & MacGillivray, H.T. 1989, MNRAS, 238,379.
 Hickson, P. 1982. ApJ, 255,382.
 Huchra, J. and Geller, M. 1982. ApJ, 257,423.
 Jerjen, H., Tammann, G.A., & Bingelli, B. 1992, Morphological and Physical Classification of Galaxies, ed. G. Longo, et al., 17.
 Lin H., Kishner R.P., Shectman S.A., Landy S.D., Oelmer A., Tucker D.L. & Schechter P.L. 1996, ApJ, 464, 60.
 López-Cruz, O. & Yee, H.K.C. 1995, ASP Conf. Ser., Vol. 86, p. 279.
 López-Cruz, O., Yee, H.K.C., Brown, J.P., Jones, C. & Forman, W. 1997, ApJ, 475,97.
 Loveday J., Peterson B.A., Efstathiou G. & Maddox S.J. 1992, ApJ, 390, 338.
 Maia, M.A.G., da Costa, L.N. and Latham, D.W. 1989. ApJS, 69,809.
 Marzke, R.O., Geller, M.J., Huchra, J.P., & Corwin, H.G. 1994, AJ, 108, 437.
 Materne, J. 1978. A&A, 63,401.
 Oemler, Jr., A. 1974, ApJ, 194, 1.
 Oegerle, W.R., Hoessel, J.G., & Ernst, R.M. 1986, AJ, 91, 697.
 Oegerle, W.R., Hoessel, J.G., & Jewison, M.S. 1987, AJ, 93, 519.
 Paturel, G., Fouque, P., Bottinelli, L. and Gouguenheim, L. 1989a. A&A Suppl. Ser. 80,299.
 Paturel, G., Fouque, P., Bottinelli, L. and Gouguenheim, L. 1989b. Extragalactic Data Base Monographs 1, vol. 1, 2 3.
 Postman, M. and Geller, M.J. 1984. ApJ, 281,95.
 Press, W.H., Flannery, B.P., Teukolsky, S.A., & Vetterling, W.T. 1987. Numerical Recipes. (Cambridge University Press, Melbourne).
 Rousseau J., Di Nella H., Paturel G., & Pettit C. MNRAS, 1996, in press.
 Schechter, P. 1976, ApJ, 203,297.
 Trentham N. 1997, MNRAS, 290,334.
 Turner, E.L. and Gott, J.R. 1976. ApJS, 32,409.
 Valotto, C.A., Nicotra, M.A., Muriel, H. and Lambas D.G. 1997. ApJ, 479,90.
 Willick, J.A., Courteau, S., Faber, S.M., Burstein, D., Dekel, A. and Kolatt, T. 1996. ApJ, 457,460.
 Willick, J.A., Courteau, S., Faber, S.M., Burstein, D., Dekel, A. and Strauss, M.A. 1997. Preprint.
 Willmer, C.N.A., Focardi, P., Chan, R., Pellegrini, P.S., & DaCosta, L.N. 1990, AJ, 101, 57.